CLIMATE MONITORING AND PREDICTION FOR SRI LANKA

HIGHLIGHTS

Rainfall Prediction

expected for the rest of the country during 19th - 23rd August.

Monitored Rainfalls



daily rainfall over Sri Lanka was 0.4 mm and hydro catchment areas have received 0.9 mm on average.

Monitored Wind



experienced at 850 mb level over the island. Upto 10m/s of south-westerlies can be expected for the next week.


Monitored Sea & Land Temp

the north of Sri Lanka.Land surface temperature remained near normal.

Monitoring


Rainfall

Daily Estimates for Rainfall from 9th August – 16th August 2022




9 August







11 August




12 August




13 August



14 August



15 August



16 August



Federation for Environment, Climate and Technology

c/o, Maintenance Office, Mahaweli Authority, Digana Village, Rajawella, Sri Lanka. Phone (+94) 81-2376746, (+94) 81-2300415

Web Site: www.fect.lk E mail: info@fect.lk LI: www.linkedin.com/in/fectlk TW: www.twitter.com/fectlk

FB: www.facebook.com/fectlk

80 100 120 140 160 180 Estimated Precipitation [mm/day]

Ocean State (Text Courtesy IRI)

Pacific sea state: August 10, 2022

Equatorial sea surface temperatures (SSTs) are below average across most of the Pacific Ocean in early-August. The tropical Pacific atmosphere is consistent with La Niña. A large majority of the models indicate, a La Niña is favored to continue through 2022 with the odds for La Niña decreasing into the Northern Hemisphere late summer (July - September 2022) before increasing through the Northern Hemisphere fall and early winter 2022.

Indian Ocean State

Sea surface temperature around Sri Lanka was above 0.5°C to the north of Sri Lanka. Across the Indian Ocean, a classical negative Indian Ocean Dipole prevails as is typical during a La Niña.

Predictions

Rainfall.

14-day prediction: NOAA NCEP models

From 17th August – 23rd August:

Total rainfall by Provinces:

Rainfall	Provinces	
55 mm	Central	
45 mm	Sabaragamuwa	
35 mm	North-central, North-western, Western, Eastern, Uva	
25 mm	Southern, Northern	

From 24th August - 30th August:

Total rainfall by Provinces:

Rainfall	Provinces
> 100 mm	Sabaragamuwa, Central, Southern, Western
95 mm	Uva, North-western
85 mm	North-central, Eastern
55 mm	Northern

MJO based OLR predictions

For the next 15 days:

MJO shall near-neutral during $17^{th}-26^{th}$ August and slightly enhance the rainfall during $27^{th}-31^{st}$ August.

Interpretation

Monitoring

Rainfall: During the last two weeks, there had been fairly heavy rainfall over the following area: Randenigala

Daily Average Rainfall in the Met stations for previous week of (9th Aug - 16th Aug) = 0.4 mm Rmax: 15.8 mm & Rmin: 0.0 mm.

Region	Average rainfall for the Last 8 days
Northern Plains	0.4 mm
Eastern	0.6 mm
Western	0.1 mm
Southern Plains	0.0 mm

The Hydro Catchment Areas recorded 0.9 mm of average rainfall for the last week

Rmax: 55.5 mm & Rmin: 0.0 mm.

Wind: South-westerly prevailed in the sea area and around the island last week.

Temperatures: The temperature anomalies were above normal for the Sabaragamuwa province, driven by the warm SST's.

Predictions_

Rainfall: During the next week (19th - 23rd August) fairly heavy rainfall is predicted for Central province and less rainfall is expected for the rest of the country.

Temperatures: The temperature remains slightly above normal in the Eastern, Northern, Northcentral and Uva provinces during 19th – 24th August.

Teleconnections:

La Niña - La Niña is favored to continue through 2022 with the odds for La Niña decreasing into the Northern Hemisphere late summer (July-September 2022) before increasing through the Northern Hemisphere fall and early winter 2022.

MJO shall near-neutral during $17^{th}-26^{th}$ August and slightly enhance the rainfall during $27^{th}-31^{st}$ August.

Seasonal Precipitation:

The precipitation forecast for the September-October-November season shows a higher tendency for below-normal precipitation for the country.

Terminology for Rainfall Ranges

	Rainfall (During 24 hours of period)
Light Showers	Less than 12.5 mm
Light to Moderate	Between 12.5 mm and 25 mm
Moderate	Between 25 mm and 50 mm
Fairly Heavy	Between 50 mm and 100 mm
Heavy	Between 100 mm and 150 mm
Very Heavy	More than 150 mm

Tropical Climate Guarantee, Federation of Environment, Climate and Technology, Columbia University Water Center, ¹ International Research Institute for Climate and Society, , Earth Institute at Columbia University, New York.





FECT Blog

Past reports available at http://fectsl.blogspot.com/



Facebook

www.facebook.com/fectlk



Twitter

www.twitter.com/fectlk

FEDERATION FOR ENVIRONMENT, CLIMATE AND TECHNOLOGY

www.climate.lk www.fect.lk

Weekly Climate Bulletin for Sri Lanka

Inside This Issue

- Monitoring
 a. Daily Rainfall Monitoring
 b. Weekly Rainfall Monitoring
 c. Monthly Rainfall Monitoring
 d. Dekadal (10 Day) Satellite Derived Rainfall Estimates
 Monitoring
 - e. Weekly Temperature Monitoring f. Weekly Wind Monitoring
- g. Weekly Average SST Anomalies
 2. Predictions


- a. NCEP GFS Ensemble 1-14 day Rainfall Predictions b. GFS (T574) Model Rainfall Forecast from RMSC New Delhi c. MJO Related OLR Forecast
- d. Weekly Temperature Forecast e. Weekly Wind Forecast
- Seasonal Predictions from IRI



MONITORING


Daily Rainfall Monitoring

The following figures show the satellite observed rainfall in the last 7 days in Sri Lanka.




Weekly Rainfall Monitoring


The following figures show the total satellite observed rainfall in the last week in Sri Lanka. The figure in the left is the total 7-day rainfall from NOAA Climate Prediction Center (CPC) Unified Precipitation Analysis and the figure in the right is the total 7-day rainfall from CPC RFE 2.0 Satellite Rainfall Estimates. The bottom two figures are the respective anomalies.




Monthly Rainfall Monitoring

The figure in the left shows the average observed rainfall in the previous month. The rainfall anomaly in the previous month is shown in the figure to the right. The brown color in the anomaly figure shows places which received less rainfall than the historical average while the green color shows places with above average rainfall. Darker shades show higher magnitudes in rainfall







The following figure shows the observed accumulated rainfall (top) and daily observed rainfall (bottom) in Sri Lanka in the last 30 days.



Dekadal (10 Day) Satellite Derived Rainfall Estimates




Weekly Temperature Monitoring




Weekly Wind Monitoring


The following figures show the mean vector wind total of the past 7 days near Sri Lanka at two levels. The figure on the left shows 850 mb (~1500 m) level and the figure on the right shows 700 mb (~3000 m) level.




Weekly average Sea Surface Temperature (SST) anomaly in the world from NOAA NCEP




Optimum Interpolated Sea Surface Temperature Anomaly in the Indian Ocean from NOAA CPC





NCEP GFS 1-14 Day prediction




IMD GFS (T574) Model Rainfall Forecast from RMSC New Delhi, India











Madden Julian Oscillation (MJO) related Outgoing Longwave Radiation (OLR) Forecast

The Outgoing Longwave Radiation (OLR) is a proxy for rainfall. This can be used to identify convective rain clouds based on the MJO phase. Violet and Blue shading indicates enhanced tropical weather and Orange shading indicates suppressed conditions. The following figure shows the forecasts of MJO associated anomolous OLR for the next 15 days from the Constructed Analogue (CA) model forecasts.

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (16 Aug 2022)




Weekly Minimum and Maximum Temperature prediction from the GFS model (from NOAA CPC)



Weekly Wind Forecast

Weekly mean vector wind total prediction from the GFS model at 850 mb (left) and 700 mb (right) levels. (from NOAA CPC)




Seasonal Rainfall and Temperature Forecast

Following is the latest seasonal precipitation and temperature prediction for the next 3 months by the IRI. The color shading indicates the probability of the most dominant tercile -- that is, the tercile having the highest forecast probability. The color bar alongside the map defines these dominant tercile probability levels. The upper side of the color bar shows the colors used for increasingly strong probabilities when the dominant tercile is the above-normal tercile, while the lower side shows likewise for the below-normal tercile. The gray color indicates an enhanced probability for the near-normal tercile (nearly always limited to 40%).

IRI Multi-Model Probability Forecast for Precipitation for September-October-November 2022, Issued August 2022

IRI Multi–Model Probability Forecast for Temperature for September–October–November 2022, Issued August 2022



Precipitation Forecast

Temperature Forecast

About us

FECT is a federation of 7 organi Zations registered in four countries which works in countries across the Indian Ocean Islands and its littoral. Over the last 20 years, we have had operations in Africa, South Asia, South-East Asia but now it is mostly in the Indian Ocean Islands.

Contact us

Federation for Environment, Climate & Technology Digana Village, Rajawella, KY20180, SRI LANKA

email: info@fect.lk phone: (+94) 81 2376746

