25 NOVEMBER 2022

CLIMATE MONITORING AND PREDICTION FOR SRI LANKA

HIGHLIGHTS

Rainfall Prediction

Moderately heavy
 (≥ 40 mm) rainfall is
 predicted for the
 Sabaragamuwa,
 Uva, and Eastern
 provinces during
 24th - 30th
 November.

Monitored Rainfalls

• During the last week, the average daily rainfall over Sri Lanka was 0.8 mm & hydro catchment areas received 0.8 mm on average and the highest average rainfall (1.6 mm) was received to the Eastern plains of the country.

Monitored Wind



- •From 14th 20th Nov, up to 6m/s of northeasterly winds were experienced at 850 mb level over the island.
- Wind Direction has changed from arriving from South-West to arriving from North East.


Monitored Sea & Land Temp

- •Sea surface temperature around Sri Lanka was above normal to the Southern half of the country.
- Land surface temperature remained near normal.


Monitoring

Rainfall _____


Daily Estimates for Rainfall from 14th November – 21st November 2022




14 November




15 November




16 November




17 November



18 November




19 November



20 November

80 100 120 140 160 180 Estimated Precipitation [mm/day]



21 November



Federation for Environment, Climate and Technology

c/o, Maintenance Office, Mahaweli Authority, Digana Village, Rajawella, Sri Lanka. Phone (+94) 81-2376746, (+94) 81-2300415

Web Site: www.fect.lk E mail: info@fect.lk LI: www.linkedin.com/in/fectlk FB: www.facebook.com/fectlk TW: www.twitter.com/fectlk

Ocean State (Text Courtesy IRI)

Pacific sea state: November 21, 2022

Equatorial sea surface temperatures (SSTs) are below average across most of the Pacific Ocean late - November. The tropical Pacific atmosphere is consistent with La Niña. A large majority of the models indicate La Niña is favored to continue through during the Northern Hemisphere winter (December-February) 2022-23, with a transition to ENSO-neutral favored in February-April 2023 (57% chance).

Indian Ocean State

Sea surface temperature around Sri Lanka was above 0.5°C to the Southern half of the country in 16th November, 2022. Across the Indian Ocean, a classical negative Indian Ocean Dipole prevails as is typical during a La Niña.

Predictions

Rainfall

7-day prediction: IMD GFS models From 24th November – 30th November:

Total rainfall by Provinces:

Rainfall	Provinces	
70 mm	Uva, Sabaragamuwa	
40 mm	Eastern	
20 mm	Southern, Central, North Western, North Central	
≤ 10 mm	Northern, Western	

MJO based **OLR** predictions

For the next 15 days:

MJO shall moderately suppress the rainfall during $23^{rd} - 27^{th}$ November, slightly suppress the rainfall during 28^{th} November -2^{nd} December, and remain near normal during $3^{rd} - 7^{th}$ December for Sri Lanka.

Interpretation

Monitoring	
0	

Rainfall: During the last two weeks, there had been very heavy rainfall over the following areas: Kegalle, Avissawella

Daily Average Rainfall in the Met stations for previous week of (15th November – 22nd November) = 0.8 mm

Rmax: 30.5 mm & Rmin: 0.0 mm.

Region	Average rainfall for the Last 8 days
Northern Plains	0.0 mm
Eastern	1.6 mm
Western	0.5 mm
Southern Plains	0.8 mm

The Hydro Catchment Areas recorded 0.8 mm of average rainfall for the last week

Rmax: 14.8 mm & Rmin: 0.0 mm.

Wind: North-easterly winds prevailed in the sea area and around the island last week.

Temperatures: The temperature anomalies were below normal for some parts of the North Central, North Eastern and Central provinces, driven by the warm SST's.

Predictions

Rainfall: During the next week $(24^{th} - 30^{th} \text{ November})$, moderately heavy ($\geq 40 \text{ mm}$) rainfall is predicted for the Uva, Sabaragamuwa, and Eastern provinces; and less rainfall is expected for the rest of the country.

Temperatures: The temperature will remain below normal for some parts of the Central and Uva provinces during $24^{th} - 30^{th}$ November.

Teleconnections: La Niña is favored to continue through during the Northern Hemisphere winter (December-February) 2022-23, with a transition to ENSO-neutral favored in February-April 2023 (57% chance).

MJO shall moderately suppress the rainfall during $23^{rd} - 27^{th}$ November, slightly suppress the rainfall during 28^{th} November -2^{nd} December, and remain near normal during $3^{rd} - 7^{th}$ December for Sri Lanka.

Seasonal Precipitation: The precipitation forecast for the December-January-February 2023 season shows a higher tendency for near-normal precipitation to the country.

Terminology for Rainfall Ranges

	Rainfall (During 24 hours of period)
Light Showers	Less than 12.5 mm
Light to Moderate	Between 12.5 mm and 25 mm
Moderate	Between 25 mm and 50 mm
Fairly Heavy	Between 50 mm and 100 mm
Heavy	Between 100 mm and 150 mm
Very Heavy	More than 150 mm

Tropical Climate Guarantee, Federation of Environment, Climate and Technology, Columbia University Water Center, ¹ International Research Institute for Climate and Society, , Earth Institute at Columbia University, New York.





FECT Blog

Past reports available at http://fectsl.blogspot.com/



Facebook



Twitter

www.facebook.com/fectlk

www.twitter.com/fectlk



FEDERATION FOR ENVIRONMENT, CLIMATE AND TECHNOLOGY

www.fect.lk www.climate.lk

Weekly Climate Bulletin for Sri Lanka

Inside This Issue

- Ionitoring
 a. Daily Rainfall Monitoring
 b. Weekly Rainfall Monitoring
 c. Monthly Rainfall Monitoring
 d. Dekadal (10 Day) Satellite Derived Rainfall Estimates
 e. Weekly Temperature Monitoring
 f. Weekly Wind Monitoring

- Weekly Average SST Anomalies

2. Predictions


- a. NCEP GFS Ensemble 1-14 day Rainfall Predictions b. GFS (T574) Model Rainfall Forecast from RMSC New Delhi c. MJO Related OLR Forecast
- Weekly Temperature Forecast Weekly Wind Forecast Seasonal Predictions from IRI



MONITORING


Daily Rainfall Monitoring

The following figures show the satellite observed rainfall in the last 7 days in Sri Lanka.




Weekly Rainfall Monitoring


The following figures show the total satellite observed rainfall in the last week in Sri Lanka. The figure in the left is the total 7-day rainfall from NOAA Climate Prediction Center (CPC) Unified Precipitation Analysis and the figure in the right is the total 7-day rainfall from CPC RFE 2.0 Satellite Rainfall Estimates. The bottom two figures are the respective anomalies.




Monthly Rainfall Monitoring

The figure in the left shows the average observed rainfall in the previous month. The rainfall anomaly in the previous month is shown in the figure to the right. The brown color in the anomaly figure shows places which received less rainfall than the historical average while the green color shows places with above average rainfall. Darker shades show higher magnitudes in rainfall







The following figure shows the observed accumulated rainfall (top) and daily observed rainfall (bottom) in Sri Lanka in the last 30 days.



(updated on 00Z22NOV2022)




Weekly Temperature Monitoring




Weekly Wind Monitoring


The following figures show the mean vector wind total of the past 7 days near Sri Lanka at two levels. The figure on the left shows 850 mb (\sim 1500 m) level and the figure on the right shows 700 mb (\sim 3000 m) level.




Weekly average Sea Surface Temperature (SST) anomaly in the world from NOAA NCEP




Optimum Interpolated Sea Surface Temperature Anomaly in the Indian Ocean from NOAA CPC





NCEP GFS 1-14 Day prediction




IMD GFS (T574) Model Rainfall Forecast from RMSC New Delhi, India











Madden Julian Oscillation (MJO) related Outgoing Longwave Radiation (OLR) Forecast


The Outgoing Longwave Radiation (OLR) is a proxy for rainfall. This can be used to identify convective rain clouds based on the MJO phase. Violet and Blue shading indicates enhanced tropical weather and Orange shading indicates suppressed conditions. The following figure shows the forecasts of MJO associated anomolous OLR for the next 15 days from the Constructed Analogue (CA) model forecasts.

OLR prediction of MJO-related anomalies using CA model reconstruction by RMM1 & RMM2 (22 Nov 2022)




Weekly Minimum and Maximum Temperature prediction from the GFS model (from NOAA CPC)






Weekly Wind Forecast

Weekly mean vector wind total prediction from the GFS model at 850 mb (left) and 700 mb (right) levels. (from NOAA CPC)






Seasonal Rainfall and Temperature Forecast

Following is the latest seasonal precipitation and temperature prediction for the next 3 months by the IRI. The color shading indicates the probability of the most dominant tercile -- that is, the tercile having the highest forecast probability. The color bar alongside the map defines these dominant tercile probability levels. The upper side of the color bar shows the colors used for increasingly strong probabilities when the dominant tercile is the above-normal tercile, while the lower side shows likewise for the below-normal tercile. The gray color indicates an enhanced probability for the near-normal tercile (nearly always limited to 40%).

IRI Multi–Model Probability Forecast for Precipitation for December–January–February 2023, Issued November 2022

IRI Multi–Model Probability Forecast for Temperature for December–January–February 2023, Issued November 2022



Precipitation Forecast

Temperature Forecast

About us

FECT is a federation of 7 organi zations registered in four countries which works in countries across the Indian Ocean Islands and its littoral. Over the last 20 years, we have had operations in Africa, South Asia, South-East Asia but now it is mostly in the Indian Ocean Islands.

Contact us

Federation for Environment, Climate & Technology 76/2 Matale Road, Akurana Kandy KY20850 SRI LANKA

email:info@fect.lk phone: (+94) 81 2376746

