CLIMATE MONITORING AND PREDICTION FOR SRI LANKA

HIGHLIGHTS

Rainfall Prediction

Heavy rainfall
 (>100mm) is
 predicted for
 Kalutara, Ratnapura
 & Galle districts;
 fairly heavy showers
 (>55 mm) will occur
 in North-western &
 central provinces
 during 29 June-5 July.

Monitored Rainfalls



• During the last week, the average daily rainfall over Sri Lanka was 1.0 mm and hydro catchment areas have received 3.5 mm on average.

Monitored Wind



•From 20th - 26th
June, up to 15 m/s
of westerly to
south-westerly
winds were
experienced over
the island.


Monitored Sea & Land Temp

•Sea surface temperature was above average of 0.5°C around Sri Lanka. Land surface temperature remained near normal.


Monitoring

Rainfall


Daily Estimates for Rainfall from 20th – 27th June 2022




20 June




21 June




22 June




23 June



24 June




25 June



26 June

80 100 120 140 160 180 Estimated Precipitation [mm/day]



27 June



Federation for Environment, Climate and Technology

c/o, Maintenance Office, Mahaweli Authority, Digana Village, Rajawella, Sri Lanka. Phone (+94) 81-2376746, (+94) 81-2300415

Web Site: www.fect.lk E mail: info@fect.lk LI: www.linkedin.com/in/fectlk FB: www.facebook.com/fectlk TW: www.twitter.com/fectlk

Ocean State (Text Courtesy IRI)

Pacific sea state: June 22, 2022

Equatorial sea surface temperatures (SSTs) are below average across most of the Pacific Ocean in late-June. The tropical Pacific atmosphere is consistent with La Niña. A large majority of the models indicate, though La Niña is favored to continue through the end of the year, the odds for La Niña decrease into the Northern Hemisphere late summer before slightly increasing through the Northern Hemisphere fall and early winter 2022.

Indian Ocean State

Sea surface temperature was above average of 0.5°C around Sri Lanka.

Predictions

Rainfall

14-day prediction: NOAA NCEP models

From 29th June – 5th July:

Total rainfall by Provinces:

Rainfall	Provinces
125 mm	Sabaragamuwa, Southern
115 mm	Western
85 mm	North-western
75 mm	Uva, Central
55 mm	Eastern, North-central
45 mm	Northern

From 6th – 12th July:

Total rainfall by Provinces:

Rainfall	Provinces
95 mm	Sabaragamuwa, Southern
85 mm	Southern
55 mm	North-western
45 mm	Uva, Central
25 mm	North Central, Eastern, Northern

MJO based OLR predictions

For the next 15 days:

MJO shall slightly enhance the rainfall during 29th June - 8th July; and near neutral during 9th - 13th July.

Interpretation

Monitoring

Rainfall: During the last two weeks, there had been fairly heavy rainfall over the following area: Galle Daily Average Rainfall in the Met stations for previous week of (20th - 27th June) = 1.0 mm Rmax: 29.6 mm & Rmin: 0.0 mm.

Region	Average rainfall for the Last 8 days
Northern Plains	0.1 mm
Eastern	0.7 mm
Western	2.5 mm
Southern Plains	0.1 mm

The Hydro Catchment Areas recorded 3.5 mm of average rainfall for the last week

Rmax: 26.5 mm & Rmin: 0 mm.

Wind: Westerly to south-westerly prevailed in the sea area surrounding the island last week.

Temperatures: The temperature anomalies were near-normal for the island, driven by the warm

SST's.

Predictions

Rainfall: During the next week (29thJune - 5thJuly) heavy rainfall (>100 mm) is predicted for the Kalutara, Ratnapura and Galle districts; fairly heavy showers (>55 mm) will occur in North-western & central provinces and the rest of the country is expected to have less rainfall.

Temperatures: The temperature remains above normal in the Northern, North-eastern, Uva and Eastern provinces during $1^{St} - 6^{th}$ July.

Teleconnections:

La Niña - Though La Niña is favored to continue through the end of the year, the odds for La Niña decrease into the Northern Hemisphere late summer (July-September2022) before slightly increasing through the Northern Hemisphere fall and early winter 2022.

MJO shall slightly enhance the rainfall during 29^{th} June - 8^{th} July; and near neutral during 9^{th} - 13^{th} July.

Seasonal Precipitation:

The precipitation forecast for the July-August-September season shows a higher tendency for abovenormal precipitation for the Jaffna district, and for the rest of the districts there is a tendency to the neutral tercile.

Terminology for Rainfall Ranges

	Rainfall (During 24 hours of period)
Light Showers	Less than 12.5 mm
Light to Moderate	Between 12.5 mm and 25 mm
Moderate	Between 25 mm and 50 mm
Fairly Heavy	Between 50 mm and 100 mm
Heavy	Between 100 mm and 150 mm
Very Heavy	More than 150 mm

Tropical Climate Guarantee, Federation of Environment, Climate and Technology, Columbia University Water Center, ¹ International Research Institute for Climate and Society, , Earth Institute at Columbia University, New York.





FECT Blog

Past reports available at http://fectsl.blogspot.com/



Facebook

www.facebook.com/fectlk



Twitter

www.twitter.com/fectlk

FEDERATION FOR ENVIRONMENT, CLIMATE AND TECHNOLOGY

www.fect.lk www.climate.lk

Weekly Climate Bulletin for Sri Lanka

Inside This Issue

- Monitoring
 a. Daily Rainfall Monitoring
 b. Weekly Rainfall Monitoring
 c. Monthly Rainfall Monitoring
 d. Dekadal (10 Day) Satellite Derived Rainfall Estimates
 Monitoring
 - e. Weekly Temperature Monitoring f. Weekly Wind Monitoring
- g. Weekly Average SST Anomalies
 2. Predictions


- a. NCEP GFS Ensemble 1-14 day Rainfall Predictions b. GFS (T574) Model Rainfall Forecast from RMSC New Delhi c. MJO Related OLR Forecast
- d. Weekly Temperature Forecast
 e. Weekly Wind Forecast
- Seasonal Predictions from IRI



MONITORING


Daily Rainfall Monitoring

The following figures show the satellite observed rainfall in the last 7 days in Sri Lanka.




Weekly Rainfall Monitoring


The following figures show the total satellite observed rainfall in the last week in Sri Lanka. The figure in the left is the total 7-day rainfall from NOAA Climate Prediction Center (CPC) Unified Precipitation Analysis and the figure in the right is the total 7-day rainfall from CPC RFE 2.0 Satellite Rainfall Estimates. The bottom two figures are the respective anomalies.




Monthly Rainfall Monitoring

The figure in the left shows the average observed rainfall in the previous month. The rainfall anomaly in the previous month is shown in the figure to the right. The brown color in the anomaly figure shows places which received less rainfall than the historical average while the green color shows places with above average rainfall. Darker shades show higher magnitudes in rainfall







The following figure shows the observed accumulated rainfall (top) and daily observed rainfall (bottom) in Sri Lanka in the last 30 days.



Dekadal (10 Day) Satellite Derived Rainfall Estimates




Weekly Temperature Monitoring




Weekly Wind Monitoring


The following figures show the mean vector wind total of the past 7 days near Sri Lanka at two levels. The figure on the left shows 850 mb (~1500 m) level and the figure on the right shows 700 mb (~3000 m) level.




Weekly average Sea Surface Temperature (SST) anomaly in the world from NOAA NCEP




Optimum Interpolated Sea Surface Temperature Anomaly in the Indian Ocean from NOAA CPC





NCEP GFS 1-14 Day prediction



IMD GFS (T574) Model Rainfall Forecast from RMSC New Delhi, India









Mad den Julian O scillation (MJO) related Outgoing Longwave Radiation (OLR) Forecast


The Outgoing Longwave Radiation (OLR) is a proxy for rainfall. This can be used to identify convective rain clouds based on the MJO phase. Violet and Blue shading indicates enhanced tropical weather and Orange shading indicates suppressed conditions. The following figure shows the forecasts of MJO associated anomolous OLR for the next 15 days from the Constructed Analogue (CA) model forecasts.



Weekly Temperature Forecast


Weekly Minimum and Maximum Temperature prediction from the GFS model (from NOAA CPC)






Weekly Wind Forecast

Weekly mean vector wind total prediction from the GFS model at 850 mb (left) and 700 mb (right) levels. (from NOAA CPC)






Seasonal Rainfall and Temperature Forecast

Following is the latest seasonal precipitation and temperature prediction for the next 3 months by the IRI. The color shading indicates the probability of the most dominant tercile -- that is, the tercile having the highest forecast probability. The color bar alongside the map defines these dominant tercile probability levels. The upper side of the color bar shows the colors used for increasingly strong probabilities when the dominant tercile is the above-normal tercile, while the lower side shows likewise for the below-normal tercile. The gray color indicates an enhanced probability for the near-normal tercile (nearly always limited to 40%).



IRI Multi–Model Probability Forecast for Temperature for July–August–September 2022, Issued June 2022



FECT is a federation of 7 organi ations registered in four countries which works in countries across the Indian Ocean Islands and its littoral. Over the last 20 years, we have had operations in Africa, South Asia, South-East Asia but now it is mostly in the Indian Ocean Islands.

Contact us

Federation for Environment, Climate & Technology Digana Village, Rajawella, KY20180, SRI LANKA

email: info@fect.lk phone: (+94) 81 2376746

