CLIMATE MONITORING AND PREDICTION FOR SRI LANKA

HIGHLIGHTS

Rainfall Prediction



Monitored Rainfalls

Up to 55 - 95 mm fairly heavy rainfall is expected in Sabaragamuwa, Western, Southern, Northern, North Central, North Western, Uva & Central provinces during 6th-10th May.

пипы



• During the last week, average daily rainfall over Sri Lanka was 6.6 mm and hydro catchment areas have received 9.9 mm on average.



•From 25th April -1st May, up to 3 m/s Southwesterlies were experienced over the Island.




Monitored Sea Surface

• Sea surface temperature was above 0.5 °C around the Island. Unlike the scorching heat in the rest of South Asia, land in Sri Lanka remained seasonable.


Monitoring

Rainfall


Daily Estimates for Rainfall from 25th April – 2nd May 2022




25 April




26 April




27 April




28 April




29 April





1 May



2 May

0 20 40 60 80 100 120 140 160 180
Estimated Precipitation [mm/day]

Federation for Environme

Federation for Environment, Climate and Technology

c/o, Maintenance Office, Mahaweli Authority, Digana Village, Rajawella, Sri Lanka. Phone (+94) 81-2376746, (+94) 81-2300415

Web Site: www.fect.lk E mail: info@fect.lk LI: www.linkedin.com/in/fectlk FB: www.facebook.com/fectlk TW: www.twitter.com/fectlk



Ocean State (Text Courtesy IRI)

Pacific sea state: April 27, 2022

Equatorial sea surface temperatures (SSTs) are below average across the most of the Pacific Ocean in late-April. The tropical Pacific atmosphere is consistent with La Niña. A large majority of the models indicate La Niña is favored to continue into the Northern Hemisphere summer.

Indian Ocean State

Sea surface temperature was above 0.5°C around the Island Unlike the scorching heat in the rest of South Asia, land in Sri Lanka remained seasonable.

Predictions

Rainfall

14-day prediction: NOAA NCEP models

From 4th – 10th May:

Total rainfall by Provinces:

Rainfall	Provinces	
95 mm	Western, Sabaragamuwa	
75 mm	Southern	
65 mm	North Western, Northern, North Central	
55 mm	Central, Uva	
45 mm	Eastern	

From 11th – 17th May:

Total rainfall by Provinces:

Rainfall	Provinces
125 mm	Western
115 mm	Sabaragamuwa
105 mm	Southern
95 mm	North Western
85 mm	Uva, Central, North Central, Northern
65 mm	Eastern

MJO based OLR predictions

For the next 15 days:

MJO shall moderately enhance the rainfall during 4th - 13th May and neutral during 14th - 18th May.

Interpretation

Monitoring -

Rainfall: During the last two weeks, there had been heavy rainfall over the following area: Kalutara Daily Average Rainfall in the Met stations for previous week of $(25^{th} \text{ April- } 2^{nd} \text{ May}) = 6.6 \text{ mm}$ Rmax: 107.2 mm & Rmin: 0.0 mm.

Region	Average rainfall for the Last 8 days
Northern Plains	4.3 mm
Eastern	6.2 mm
Western	10 mm
Southern Plains	3.3 mm

The Hydro Catchment Areas recorded 9.9 mm of average rainfall for the last week

Rmax: 98 mm & Rmin: 0 mm.

Wind: South-westerly prevailed in the sea area surrounding the island last week.

Temperatures: The temperature anomalies were near neutral for the country, driven by the warm SST's.

Predictions

Rainfall: During the next week (6th - 10th May) fairly heavy rainfall is predicted for the Western, Sabaragamuwa, Southern, Northern, North Central, North Western, Uva and Central provinces.

Temperatures: The temperature remains slightly below normal in the central provinces and above normal in the eastern, uva and northern province during 6th - 14th May.

Teleconnections:

La Nina - The SST forecast indicates that La Niña is favored to continue into the Northern Hemisphere summer (June-August 2022).

MJO shall moderately enhance the rainfall during 4^{th} - 13^{th} May and neutral during 14^{th} – 18^{th} May.

Seasonal Precipitation:

The precipitation forecast for the May-June-July season shows below-normal precipitation for the island, but above-normal precipitation for the northern province.

Terminology for Rainfall Ranges

	Rainfall (During 24 hours of period)
Light Showers	Less than 12.5 mm
Light to Moderate	Between 12.5 mm and 25 mm
Moderate	Between 25 mm and 50 mm
Fairly Heavy	Between 50 mm and 100 mm
Heavy	Between 100 mm and 150 mm
Very Heavy	More than 150 mm

Tropical Climate Guarantee, Federation of Environment, Climate and Technology, Columbia University Water Center, ¹ International Research Institute for Climate and Society, , Earth Institute at Columbia University, New York.









FEDERATION FOR ENVIRONMENT, CLIMATE AND TECHNOLOGY

www.fect.lk www.climate.lk

Weekly Climate Bulletin for Sri Lanka

Inside This Issue

- Monitoring
 a. Daily Rainfall Monitoring
 b. Weekly Rainfall Monitoring
 c. Monthly Rainfall Monitoring
 d. Dekadal (10 Day) Satellite Derived Rainfall Estimates
 Weekly Temporature Monitoring
- e. Weekly Temperature Monitoring
 f. Weekly Wind Monitoring
 g. Weekly Average SST Anomalies


 2. Predictions


- a. NCEP GFS Ensemble 1-14 day Rainfall Predictions b. GFS (T574) Model Rainfall Forecast from RMSC New Delhi c. MJO Related OLR Forecast
- d. Weekly Temperature Forecast
 e. Weekly Wind Forecast
- Seasonal Predictions from IRI


MONITORING


Daily Rainfall Monitoring


The following figures show the satellite observed rainfall in the last 7 days in Sri Lanka.






















Weekly Rainfall Monitoring


The following figures show the total satellite observed rainfall in the last week in Sri Lanka. The figure in the left is the total 7-day rainfall from NOAA Climate Prediction Center (CPC) Unified Precipitation Analysis and the figure in the right is the total 7-day rainfall from CPC RFE 2.0 Satellite Rainfall Estimates. The bottom two figures are the respective anomalies




Monthly Rainfall Monitoring

The figure in the left shows the average observed rainfall in the previous month. The rainfall anomaly in the previous month is shown in the figure to the right. The brown color in the anomaly figure shows places which received less rainfall than the historical average while the green color shows places with above average rainfall. Darker shades show higher magnitudes in rainfall







The following figure shows the observed accumulated rainfall (top) and daily observed rainfall (bottom) in Sri Lanka in the last 30 days.



Dekadal (10 Day) Satellite Derived Rainfall Estimates




Weekly Temperature Monitoring




Weekly Wind Monitoring


The following figures show the mean vector wind total of the past 7 days near Sri Lanka at two levels. The figure on the left shows 850 mb (~1500 m) level and the figure on the right shows 700 mb (~3000 m) level.




Weekly average Sea Surface Temperature (SST) anomaly in the world from NOAA NCEP




Optimum Interpolated Sea Surface Temperature Anomaly in the Indian Ocean from NOAA CPC





NCEP GFS 1-14 Day prediction



IMD GFS (T574) Model Rainfall Forecast from RMSC New Delhi, India









Madden Julian Oscillation (MJO) related Outgoing Longwave Radiation (OLR) Forecast


The Outgoing Longwave Radiation (OLR) is a proxy for rainfall. This can be used to identify convective rain clouds based on the MJO phase. Violet and Blue shading indicates enhanced tropical weather and Orange shading indicates suppressed conditions. The following figure shows the forecasts of MJO associated anomolous OLR for the next 15 days from the Constructed Analogue (CA) model forecasts.






Weekly Minimum and Maximum Temperature prediction from the GFS model (from NOAA CPC)






Weekly Wind Forecast

Weekly mean vector wind total prediction from the GFS model at 850 mb (left) and 700 mb (right) levels. (from NOAA CPC)






Seasonal Rainfall and Temperature Forecast

Following is the latest seasonal precipitation and temperature prediction for the next 3 months by the IRI. The color shading indicates the probability of the most dominant tercile — that is, the tercile having the highest forecast probability. The color bar alongside the map defines these dominant tercile probability levels. The upper side of the color bar shows the colors used for increasingly strong probabilities when the dominant tercile is the above-normal tercile, while the lower side shows likewise for the below-normal tercile. The gray color indicates an enhanced probability for the near-normal tercile (nearly always limited to 40%).

IRI Multi–Model Probability Forecast for Precipitation for May–June–July 2022, Issued April 2022

IRI Multi–Model Probability Forecast for Temperature for May–June–July 2022, Issued April 2022



Precipitation Forecast

Temperature Forecast

About us

FECT is a federation of 7 organizations registered in four countries which works in countries across the Indian Ocean Islands and its littoral. Over the last 20 years, we have had operations in Africa, South Asia, South-East Asia but now it is mostly in the Indian Ocean Islands.

Contact us

Federation for Environment, Climate & Technology Digana Village, Rajawella, KY20180, SRI LANKA

email: info@fect.lk phone: (+94) 81 2376746

